

The Price of Anarchy in Network Creation Games

Erik D. Demaine*
Computer Science & AI Lab
M.I.T.
Cambridge, MA
edemaine@mit.edu

MohammadTaghi Hajiaghayi*
Computer Science & AI Lab
M.I.T.
Cambridge, MA
hajiagha@mit.edu

Hamid Mahini
Dept. of Computer Engineering
Sharif University of Technology
Tehran, Iran
mahini@ce.sharif.edu

Morteza Zadimoghaddam
Dept. of Computer Engineering
Sharif University of Technology
Tehran, Iran
zadimoghaddam@ce.sharif.edu

ABSTRACT

We study Nash equilibria in the setting of network creation games introduced recently by Fabrikant, Luthra, Maneva, Papadimitriou and Shenker. In this game we have a set of selfish node players, each creating some incident links, and the goal is to minimize α times the cost of the created links plus sum of the distances to all other players. Fabrikant et al. proved an upper bound $O(\sqrt{\alpha})$ on the price of anarchy, i.e., the relative cost of the lack of coordination. Albers, Eilts, Even-Dar, Mansour, and Roditty show that the price of anarchy is constant for $\alpha = O(\sqrt{n})$ and for $\alpha \geq 12n \lceil \lg n \rceil$, and that the price of anarchy is $15 \left(1 + (\min\{\frac{\alpha^2}{n}, \frac{n^2}{\alpha}\})^{1/3}\right)$ for any α . The latter bound shows the first sublinear worst-case bound, $O(n^{1/3})$, for all α . But no better bound is known for α between $\omega(\sqrt{n})$ and $o(n \lg n)$. Yet $\alpha \approx n$ is perhaps the most interesting range, for it corresponds to considering the average distance (instead of the sum of distances) to other nodes to be roughly on par with link creation (effectively dividing α by n).

In this paper, we prove the first $o(n^\epsilon)$ upper bound for general α , namely $2^{O(\sqrt{\lg n})}$. We also prove a constant upper bound for $\alpha = O(n^{1-\epsilon})$ for any fixed $\epsilon > 0$, substantially reducing the range of α for which constant bounds have not been obtained. Along the way, we also improve the constant upper bound by Albers et al. (with the lead constant of 15) to 6 for $\alpha < (n/2)^{1/2}$ and to 4 for $\alpha < (n/2)^{1/3}$.

Next we consider the bilateral network variant of Corbo and Parkes in which links can be created only with the consent of both endpoints and the link price is shared equally by the two. Corbo and Parkes show an upper bound of $O(\sqrt{\alpha})$

*Supported in part by NSF under grant number ITR ANI-0205445 and by MADALGO — Center for Massive Data Algorithmics — a Center of the Danish National Research Foundation.

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. To copy otherwise, to republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee.

PODC'07, August 12–15, 2007, Portland, Oregon, USA.
Copyright 2007 ACM 978-1-59593-616-5/07/0008 ...\$5.00.

and a lower bound of $\Omega(\lg \alpha)$ for $\alpha \leq n$. In this paper, we show that in fact the upper bound $O(\sqrt{\alpha})$ is tight for $\alpha \leq n$, by proving a matching lower bound of $\Omega(\sqrt{\alpha})$. For $\alpha > n$, we prove that the price of anarchy is $\Theta(n/\sqrt{\alpha})$.

Finally we introduce a variant of both network creation games, in which each player desires to minimize α times the cost of its created links plus the maximum distance (instead of the sum of distances) to the other players. This variant of the problem is naturally motivated by considering the worst case instead of the average case. Interestingly, for the original (unilateral) game, we show that the price of anarchy is at most 2 for $\alpha \geq n$, $O(\min\{4^{\sqrt{\lg n}}, (n/\alpha)^{1/3}\})$ for $2\sqrt{\lg n} \leq \alpha \leq n$, and $O(n^{2/\alpha})$ for $\alpha < 2\sqrt{\lg n}$. For the bilateral game, we prove matching upper and lower bounds of $\Theta(\frac{n}{\alpha+1})$ for $\alpha \leq n$, and an upper bound of 2 for $\alpha > n$.

Categories and Subject Descriptors

F.2.2 [Theory of Computation]: Analysis of Algorithms and Problem Complexity—*Nonnumerical Algorithms and Problems*; G.2.2 [Mathematics of Computing]: Discrete Mathematics—*graph theory, network problems*

General Terms

Performance, Design, Economics

Keywords

network design, routing, price of anarchy, Nash equilibrium

1. INTRODUCTION

Network design is an fundamental family of problems at the intersection between computer science and operations research, amplified in importance by the sustained growth of computer networks such as the Internet. Traditionally, the goal is to find a minimum-cost (sub) network that satisfies some specified property such as k -connectivity or connectivity on terminals (as in the classic Steiner tree problem). Such a formulation captures the (possibly incremental) creation cost of the network, but does not incorporate the cost of actually using the network. By contrast, *network routing* has the goal of optimizing the usage cost of the network, but assumes that the network has already been created.

The *network creation game* attempts to unify the network design and network routing problems by modeling both creation and usage costs. Specifically, each node in the system can create a link (edge) to any other node, at a cost of α . In addition to these creation costs, each node incurs a usage cost related to the distances to the other nodes. In the model introduced by Fabrikant, Luthra, Maneva, Papadimitriou, and Shenker at PODC 2003 [8], the usage cost incurred by a node is the sum of distances to all other nodes. Equivalently, if we divide the cost (and thus α) by the number n of nodes, the usage cost is the average distance to other nodes. In another natural model, the usage cost incurred by a node is the maximum distance to all other nodes: this model captures the worst-case instead of average-case behavior of routing.

To model the dominant behavior of large-scale networking scenarios such as the Internet, we consider each node to be an agent (player) [9] that selfishly tries to minimize its own creation and usage cost [8, 1, 5]. In this context, the *price of anarchy* [10, 13, 15] is the worst possible ratio of the total cost found by some independent selfish behavior and the optimal total cost possible by a centralized, social welfare maximizing solution. The price of anarchy is a well-studied concept in algorithmic game theory for problems such as load balancing, routing, and network design; see, e.g., [13, 6, 14, 8, 3, 2, 4, 5, 1].

Previous work.

Three papers consider the price of anarchy in the sum network creation game. Fabrikant et al. [8] prove an upper bound of $O(\sqrt{\alpha})$ on the price of anarchy for all α . Lin [12] prove that the price of anarchy is constant for two ranges of α : $\alpha = O(\sqrt{n})$ and $\alpha \geq cn^{3/2}$ for some $c > 0$. Independently, Albers et al. [1] prove that the price of anarchy is constant for $\alpha = O(\sqrt{n})$, as well as for the larger range $\alpha \geq 12n \lceil \lg n \rceil$. In addition, Albers et al. prove a general upper bound of $15 \left(1 + (\min\{\frac{\alpha^2}{n}, \frac{n^2}{\alpha}\})^{1/3}\right)$. The latter bound shows the first sublinear worst-case bound, $O(n^{1/3})$, for all α . But no better bound is known for α between $\omega(\sqrt{n})$ and $o(n \lg n)$. Yet $\alpha \approx n$ is perhaps the most interesting range, for it corresponds to considering the average distance (instead of the sum of distances) to other nodes to be roughly on par with link creation (effectively dividing α by n).

The *bilateral* variation on the network creation game, considered by Corbo and Parkes in PODC 2005 [5], requires both nodes to agree before they can create a link between them. In the sum bilateral network creation game, Corbo and Parkes prove that the price of anarchy is between $\Omega(\lg \alpha)$ and $O(\sqrt{\alpha})$ for $\alpha \leq n$. Although not claimed in their paper, the proof of their upper bound also establishes an upper bound of $O(n/\sqrt{\alpha})$ for $\alpha \geq n$.

Our results.

In the sum (unilateral) network creation game of Fabrikant et al. [8], we prove the first $o(n^\varepsilon)$ upper bound for general α , namely $2^{O(\sqrt{\lg n})}$. We also prove a constant upper bound for $\alpha = O(n^{1-\varepsilon})$ for any fixed $\varepsilon > 0$, substantially reducing the range of α for which constant bounds have not been obtained. Along the way, we also improve the constant upper bound by Albers et al. [1] (with the lead constant of 15) to 6 for $\alpha < (n/2)^{1/2}$ and to 4 for $\alpha < (n/2)^{1/3}$.

In the max (unilateral) version, we prove that the price of anarchy is at most 2 for $\alpha \geq n$, $O(\min\{4^{\sqrt{\lg n}}, (n/\alpha)^{1/3}\})$ for $2\sqrt{\lg n} \leq \alpha \leq n$, and $O(n^{2/\alpha})$ for $\alpha < 2\sqrt{\lg n}$.

Our primary proof technique can be most closely viewed as a kind of “region growing” from approximation algorithms; see, e.g., [11].

In the (sum) bilateral network creation game of Corbo and Parkes [5], we show that their $O(\min\{\sqrt{\alpha}, n/\sqrt{\alpha}\})$ upper bound is tight by proving a matching lower bound of $\Omega(\min\{\sqrt{\alpha}, n/\sqrt{\alpha}\})$. In the max (bilateral) version, we prove matching upper and lower bounds of $\Theta(\frac{n}{\alpha+1})$ for $\alpha \leq n$, and an upper bound of 2 for $\alpha > n$.

Table 1 summarizes the best bounds known for each of the three models and for all ranges of α . Interestingly, these results reveal that the price of anarchy can depend significantly on α .

2. MODELS

Formally, we define four games depending on the objective (sum or max) and the consent (unilateral or bilateral). In all versions, we have n players; call them $1, 2, \dots, n$. The strategy of player i is specified by a subset s_i of $\{1, 2, \dots, n\} - \{i\}$, which corresponds to the set of neighbors to which player i forms a link. Together, let $s = \langle s_1, s_2, \dots, s_n \rangle$ denote the strategies of all players.

To define the cost of a strategy, we introduce an undirected graph G_s with vertex set $\{1, 2, \dots, n\}$. In the unilateral game, G_s has an edge $\{i, j\}$ if either $i \in s_j$ or $j \in s_i$. In the bilateral game, G_s has an edge $\{i, j\}$ if both $i \in s_j$ and $j \in s_i$. Define $d_s(i, j)$ to be the distance between vertices i and j in graph G_s . In the sum game, the cost incurred by player i is

$$c_i(s) = \alpha |s_i| + \sum_{j=1}^n d_s(i, j),$$

and in the max game, the cost incurred by player i is

$$c_i(s) = \alpha |s_i| + \max_{j=1}^n d_s(i, j).$$

In both cases, the total cost incurred by strategy s is $c(s) = \sum_{i=1}^n c_i(s)$.

In the unilateral game, a (pure) *Nash equilibrium* is a strategy s such that $c(s) \leq c(s')$ for all strategies s' that differ from s in only one player i . The *price of anarchy* is then the maximum cost of a Nash equilibrium divided by the minimum cost of any strategy (called the *social optimum*).

In the bilateral game, Nash equilibria are not so interesting because the game requires coalition between two players to create an edge (in general). For example, if every player i chooses the empty strategy $s_i = \emptyset$, then we obtain a Nash equilibrium inducing an empty graph G_s , which has an infinite cost $c(s)$. To address this issue, Corbo and Parkes [5] define a strategy to be *pairwise stable* if (1) for any edge $\{i, j\}$ of G_s , both $c_i(s) \leq c_i(s')$ and $c_j(s) \leq c_j(s')$ where s' differs from s only in deleting edge $\{i, j\}$ from G_s ; and (2) for any nonedge $\{i, j\}$ of G_s , either $c_i(s) < c_i(s')$ or $c_j(s) < c_j(s')$ where s' differs from s only in adding edge $\{i, j\}$ to G_s . The *price of anarchy* is then the maximum cost of a pairwise-stable strategy divided by the social optimum (the minimum cost of any strategy).

We spend the bulk of this paper (Sections 3–6) on the original version of the network creation game, sum unilateral. Then we consider the max unilateral game in Section 7,

$\alpha =$	0	1	2	$2\sqrt{\lg n}$	$\sqrt[3]{n/2}$	$\sqrt{n/2}$	$O(n^{1-\varepsilon})$	n	$12n\lceil \lg n \rceil$	n^2	
Sum unilateral	1	$\frac{4}{3}$ [8]	≤ 4 (§4)	≤ 6 (§4)	$\Theta(1)$ (§5)	$2^{O(\sqrt{\lg n})}$ (§6)	$\Theta(1)$ [1]				
Max unilateral		$O(n^{2/\alpha})$ (§7)		$O(\min\{4^{\sqrt{\lg n}}, (n/\alpha)^{1/3}\})$ (§7)			≤ 2 (§7)				
Sum bilateral	1	$\Theta(\sqrt{\alpha})$ (O in [5], Ω in §8)						$\Theta(\frac{n}{\sqrt{\alpha}})$ (O in [5] ^a , Ω in §8)			
Max bilateral		$\Theta(\frac{n}{\alpha+1})$ (§9)						≤ 2 (§9)			

Table 1: Summary of best known results for all three models of network creation games and for all ranges of α . ^a[5, Proposition 4] claims $O(\sqrt{\alpha})$ for $\alpha < n^2$ but in fact proves $O(\min\{\sqrt{\alpha}, n/\sqrt{\alpha}\})$.

the sum bilateral game in Section 8, and the max bilateral game in Section 9. Thus we default to the unilateral game, often omitting the term, and within the unilateral game, we default to the sum version.

3. PRELIMINARIES

In this section, we define some helpful notation and prove some basic results. Call a graph G_s corresponding to a Nash equilibrium s an *equilibrium graph*. In such a graph, let $d_s(u, v)$ be the length of the shortest path from u to v . Let $N_k(u)$ denote the set of vertices with distance at most k from vertex u , and let $N_k = \min_{v \in G} |N_k(v)|$.

We start with a lemma proved in [1] about the sum unilateral game:

LEMMA 1. [1, proof of Theorem 3.2] *For any Nash equilibrium s and any vertex v_0 in G_s , the cost $c(s)$ is at most $2\alpha(n-1) + n \text{Dist}(v_0) + (n-1)^2$ where $\text{Dist}(v_0) = \sum_{v \in V(G_s)} d_s(v_0, v)$.*

Now we use this lemma to relate the price of anarchy to depth in a breadth-first search (BFS) tree, a connection also used in [1]. We assume henceforth that $\alpha \geq 2$, because otherwise the price of anarchy for the sum unilateral game is already known from [8]. In this case, it is known that the social optimum is attained by a star graph [8].

LEMMA 2. *If the BFS tree in a equilibrium graph G_s rooted at vertex u has depth d , then price of anarchy is at most $d + 1$.*

PROOF. By Lemma 1 and because the cost of the social optimum (a star) is at least $\alpha(n-1) + n(n-1)$, the price of anarchy is at most $\frac{2\alpha(n-1) + n \text{Dist}(u) + (n-1)^2}{\alpha(n-1) + n(n-1)}$. Because the height of the BFS tree rooted at u is d , we have $\text{Dist}(u) \leq (n-1)d$. Hence the price of anarchy is at most $\frac{2\alpha(n-1) + n(n-1)d + (n-1)^2}{\alpha(n-1) + n(n-1)} < \frac{2\alpha(n-1) + n(n-1)(d+1)}{\alpha(n-1) + n(n-1)} \leq \max\{\frac{2\alpha(n-1)}{\alpha(n-1)}, \frac{n(n-1)(d+1)}{n(n-1)}\} = \max\{2, d+1\} = d+1$. \square

LEMMA 3. *For any equilibrium graph G_s , $|N_2(u)| > n/(2\alpha)$ for every vertex u and $\alpha \geq 1$.*

PROOF. We can assume that the number vertices with distance more than 2 from u is at least $n/2$; otherwise $|N_2(u)| > n/2 \geq n/(2\alpha)$. Let S be the vertices whose distance is exactly 2 from u . For each vertex v with $d_s(u, v) \geq 2$, we pick any one of its shortest paths to u and assign v to the only vertex in this path that is in S . The number of vertices assigned to a vertex $w \in S$ can be no more than α , because otherwise vertex u could buy edge $\{u, w\}$ and decrease its distance to each vertex assigned to w by at least 1. There is one assignment for each vertex with distance at least 2 from u , so the total number of assignments to vertices in S is more than $n/2$. Therefore, $|S| > (n/2)/\alpha = n/(2\alpha)$. \square

4. IMPROVED CONSTANT UPPER BOUNDS FOR $\alpha = O(\sqrt{N})$

In this section, we use the basic results from the previous section to improve the $O(1)$ bounds for the sum unilateral game with $\alpha = O(\sqrt{n})$ obtained by Albers et al. [1]. Recall that their bound has a lead constant of (and is thus at least) 15. We prove an upper bound of 6 for $\alpha < (n/2)^{1/2}$ and 4 for $\alpha < (n/2)^{1/3}$.

THEOREM 1. *For $\alpha < (n/2)^{1/2}$, the price of anarchy is at most 6.*

PROOF. We prove that the height of the BFS tree rooted at an arbitrary vertex u in an equilibrium graph is at most 5. Suppose for contradiction that there is a vertex v at distance at least 6 from u . Vertex v can buy $\{u, v\}$ to decrease its distance from all vertices in $N_2(u)$ by at least 1. Because vertex v has not bought the edge $\{u, v\}$, we conclude that $|N_2(u)| < \alpha$. By Lemma 3, $|N_2(u)| > n/(2\alpha)$. Hence $\alpha > n/(2\alpha)$, which contradicts the hypothesis of the lemma. Therefore the height of the BFS tree rooted at u is at most 5. By Lemma 2, the price of anarchy is at most $5 + 1 = 6$. \square

THEOREM 2. *For $\alpha < (n/2)^{1/3}$, the price of anarchy is at most 4.*

PROOF. We prove that there is a choice of root vertex v in an equilibrium graph such that the height of the BFS tree is at most 3. Let Δ be the maximum degree of a vertex in the graph and let u be an arbitrary vertex. Certainly $|N_2(u)| \leq 1 + \Delta^2$. On the other hand, by Lemma 3, $|N_2(u)| > n/(2\alpha)$. By the hypothesis of the lemma, $1 + \Delta^2 > n/(2\alpha) > \alpha^2$. Hence $\Delta > \alpha - 1$. Let v be a vertex with degree equal to Δ . Suppose for contradiction that there is vertex u at distance at least 4 from v . Vertex u can buy edge $\{u, v\}$ to decrease its distance to all vertices in $N_1(v)$ by at least 1 and thus decrease its total cost by at least $|N_1(v)| = \Delta + 1 > \alpha$, contradicting equilibrium. Therefore the height of the BFS tree rooted at v is at most 3. By Lemma 2, the price of anarchy is at most $3 + 1 = 4$. \square

5. CONSTANT UPPER BOUND FOR $\alpha = O(N^{1-\varepsilon})$

In this section, we extend the range for which we know a constant upper bound on the price of anarchy in the sum unilateral game. Specifically, for $\alpha = O(n^{1-\varepsilon})$, we prove an upper bound of $2^{O(1/\varepsilon)}$. We start with some lemmas that will be useful in this section and the next.

LEMMA 4. *For any vertex u in an equilibrium graph G_s , if $|N_k(u)| > n/2$, then $|N_{2k+2\alpha/n}(u)| \geq n$.*

PROOF. We prove the contrapositive. Suppose $|N_{2k+2\alpha/n}(u)| < n$. Then there is a vertex v

with $d_s(u, v) \geq 2k + 1 + 2\alpha/n$. For every vertex $x \in N_k(u)$, $d_s(u, x) \leq k$. By the triangle inequality $d_s(u, x) + d_s(x, v) \geq d_s(u, v)$, we have $d_s(x, v) \geq k + 1 + 2\alpha/n$. If vertex v bought the edge $\{v, u\}$, then the distance between v and x would decrease by at least $2\alpha/n$, so $\text{Dist}(v)$ would decrease by at least $N_k(u) \cdot 2\alpha/n$. Because v has not bought the edge $\{v, u\}$, we have $\alpha \geq |N_k(u)| \cdot 2\alpha/n$, i.e., $|N_k(u)| \leq n/2$. \square

By setting $\alpha = n/2$ and $\alpha = 12n \lg n$, we conclude the following corollaries, which we will use in Theorems 3 and 4.

COROLLARY 1. *For any vertex u in an equilibrium graph G_s with $\alpha < n/2$, if $|N_k(u)| > n/2$, then $|N_{2k+1}(u)| \geq n$.*

COROLLARY 2. *For any vertex u in an equilibrium graph G_s with $\alpha < 12n \lg n$, if $|N_k(u)| > n/2$, then $|N_{2k+24 \lg n}(u)| \geq n$.*

LEMMA 5. *If $|N_k(u)| > Y$ for every vertex u in an equilibrium graph G_s , then either $|N_{2k+3}(u)| > n/2$ for some vertex u or $|N_{3k+3}(u)| > Yn/\alpha$ for every vertex u .*

PROOF. If there is a vertex u with $|N_{2k+3}(u)| > n/2$, then the claim is obvious. Otherwise, for every vertex u , $|N_{2k+3}(u)| \leq n/2$. Let u be an arbitrary vertex. Let S be the set of vertices whose distance from u is $2k + 3$. We select a subset of S , called *center points*, by the following greedy algorithm. First we unmark all vertices in S . Then we repeatedly select an unmarked vertex $x \in S$ as a center point, mark all unmarked vertices in S whose distance from x is at most $2k$, and assign these vertices to x .

Suppose that we select l vertices x_1, x_2, \dots, x_l as center points. We prove that $l \geq n/\alpha$. Let C_i be the vertices in S assigned to x_i . By construction, $S = \bigcup_{i=1}^l C_i$. We also assign each vertex v at distance at least $2k + 3$ from u to one of these center points, as follows. Pick any one shortest path from v to u , which contains exactly one vertex $w \in S$, and assign v to the same center point as w . Let T_i be the set of vertices assigned to x_i and whose distance from u is more than $2k + 3$. By construction, $\bigcup_{i=1}^l T_i$ is the set of vertices at distance more than $2k + 3$ from u . The shortest path from $v \in T_i$ to u uses some vertex $w \in C_i$. If u bought the edge $\{u, x_i\}$, then the distance between u and w would become at most $2k + 1$. Because $d_s(u, w) = 2k + 3$ in the current graph, buying edge $\{u, x_i\}$ would decrease u 's distance to v by at least $2k + 3 - (1 + 2k) = 2$. Because u has not bought the edge $\{u, x_i\}$, we conclude that $\alpha \geq 2|T_i|$. On the other hand, $|N_{2k+3}(u)| \leq n/2$, so $\sum_{i=1}^l |T_i| \geq n/2$. Therefore, $l\alpha \geq 2 \sum_{i=1}^l |T_i| \geq n$ and hence $l \geq n/\alpha$.

According to the greedy algorithm, the distance between any pair of center points is more than $2k$; hence, $N_k(x_i) \cap N_k(x_j) = \emptyset$ for $i \neq j$. By the hypothesis of the lemma, $|N_k(x_i)| > Y$ for every vertex x_i ; hence $|\bigcup_{i=1}^l N_k(x_i)| = \sum_{i=1}^l |N_k(x_i)| > lY$. For every $i \leq l$, we have $d_s(u, x_i) = 2k + 3$, so vertex u has a path of length at most $3k + 3$ to every vertex whose distance to x_i is at most k . Therefore, $|N_{3k+3}(u)| \geq |\bigcup_{i=1}^l N_k(x_i)| > lY \geq Yn/\alpha$. \square

THEOREM 3. *For $\varepsilon \geq 1/\lg n$ and $1 \leq \alpha < n^{1-\varepsilon}$, the price of anarchy is at most $4.667 \cdot 3^{\lceil 1/\varepsilon \rceil} + 8$.*

PROOF. Consider an equilibrium graph G_s . Let $X = n/\alpha > n^\varepsilon$. Define $a_1 = 2$ and $a_i = 3a_{i-1} + 3$ for all $i > 1$.

By Lemma 3, $N_2(u) > n/(2\alpha) = X/2$. By Lemma 5, for each $i \geq 1$, either $N_{2a_i+3}(v) > n/2$ for some vertex v or $|N_{a_i+1}(v)| > (n/\alpha)N_{a_i} = XN_{a_i}$. Let j be the least number for which $|N_{2a_j+3}(v)| > n/2$ for some vertex v . By this definition, for each $i < j$, $|N_{a_i+1}(v)| > (n/\alpha)N_{a_i} = XN_{a_i}$. Because $N_{a_1} = N_2 > X/2$, we obtain that $N_{a_i} > X^i/2$ for every $i \leq j$. On the other hand, $X^j/2 < N_{a_j} \leq n$, so $X^j < 2n$. Thus $j \leq \lceil 1/\varepsilon \rceil$, so $|N_{a_{2^{\lceil 1/\varepsilon \rceil}+3}}(v)| \geq |N_{2a_j+3}(v)| > n/2$. By Corollary 1, $|N_{4a_{\lceil 1/\varepsilon \rceil}+7}(v)| = n$. Hence, the height of the BFS tree rooted at vertex v is at most $4a_{\lceil 1/\varepsilon \rceil} + 7$. By Lemma 2, the price of anarchy is at most $4a_{\lceil 1/\varepsilon \rceil} + 8$. Solving the recurrence relation $a_i = 3a_{i-1} + 3$ with $a_1 = 2$, we obtain that $a_i = \frac{7}{6}3^i - \frac{3}{2} < \frac{7}{6}3^i$. Therefore the price of anarchy is at most $4 \frac{7}{6}3^{\lceil 1/\varepsilon \rceil} + 8 \leq 4.667 \cdot 3^{\lceil 1/\varepsilon \rceil} + 8$. \square

6. $O(N^\varepsilon)$ UPPER BOUND FOR $\alpha < 12N \lg N$

In this section, we prove the first $o(n^\varepsilon)$ bound for the sum unilateral game with α between $\Omega(n)$ and $o(n \lg n)$. Specifically, we show an upper bound of $2^{O(\sqrt{\lg n})}$. First we need the following lemma.

LEMMA 6. *If $|N_k(u)| > Y$ for every vertex u in an equilibrium graph G_s , then either $|N_{4k+1}(u)| > n/2$ for some vertex u or $|N_{5k+1}(u)| > Ykn/\alpha$ for every vertex u .*

PROOF. The proof is similar to the proof of Lemma 5. If there is a vertex u with $|N_{4k+1}(u)| > n/2$, then the claim is obvious. Otherwise, for every vertex u , $|N_{4k+1}(u)| \leq n/2$. Let u be an arbitrary vertex. Let S be the set of vertices whose distance from u is $4k + 1$. We select a subset of S , called *center points*, by the following greedy algorithm. First we unmark all vertices in S . Then we repeatedly select an unmarked vertex $x \in S$ as a center point, mark all unmarked vertices in S whose distance is from x at most $2k$, and assign these vertices to x .

Suppose that we select l vertices x_1, x_2, \dots, x_l as center points. We prove that $l \geq kn/\alpha$. Let C_i be the vertices in S assigned to x_i . By construction, $S = \bigcup_{i=1}^l C_i$. We also assign each vertex v with at distance at least $4k + 1$ from u to one of these center points. Pick any one shortest path from v to u , which contains exactly one vertex $w \in S$, and assign v to the same center point as w . Let T_i be the set of vertices assigned to x_i and whose distance from u is more than $4k + 1$. By construction, $\bigcup_{i=1}^l T_i$ is the set of vertices at distance more than $4k + 1$ from u . The shortest path from $v \in T_i$ to u uses some vertex $w \in C_i$. If u bought the edge $\{u, x_i\}$, then the distance between u and w would become at most $2k + 1$. Because $d_s(u, w) = 4k + 1$, buying edge $\{u, x_i\}$ would decrease u 's distance to v by at least $4k + 1 - (2k + 1) = 2k$. Because u has not bought the edge $\{u, x_i\}$, we conclude that $\alpha \geq 2k|T_i|$. On the other hand, $|N_{4k+1}(u)| \leq n/2$ and $\sum_{i=1}^l |T_i| \geq n/2$. Therefore, $l\alpha \geq 2k \sum_{i=1}^l |T_i| \geq kn$ and hence $l \geq kn/\alpha$.

According to the greedy algorithm, the distance between any pair of center points is more than $2k$; hence, $N_k(x_i) \cap N_k(x_j) = \emptyset$ for $i \neq j$. By the hypothesis of the lemma, $|N_k(x_i)| > Y$ for every vertex x_i ; hence $|\bigcup_{i=1}^l N_k(x_i)| = \sum_{i=1}^l |N_k(x_i)| > lY$. For every $i \leq l$, we have $d_s(u, x_i) = 4k + 1$, so vertex u has a path of length at most $5k + 1$ to every vertex whose distance to x_i is at most k . Therefore, $|N_{5k+1}(u)| \geq |\bigcup_{i=1}^l N_k(x_i)| > lY \geq Ykn/\alpha$. \square

THEOREM 4. For $1 \leq \alpha < 12n \lg n$, the price of anarchy is $O(5^{\sqrt{\lg n}} \lg n)$.

PROOF. The proof is similar to the proof of Theorem 3. Let $Z = 12 \lg n$. By the hypothesis of the theorem, $\alpha/n < 12 \lg n = Z$. Because any equilibrium graph G_s is connected, $|N_Z| > Z$. By Lemma 6, either $|N_{4k+1}(v)| > n/2$ for some vertex v or $N_{5k+1} \geq (n/\alpha)kN_k$. Define a_0, a_1, \dots by the recurrence relation $a_i = 5a_{i-1} + 1$ with $a_0 = Z$. By induction, $a_i > Z5^i$. Suppose j is the least number for which $|N_{4a_j+1}(v)| > n/2$. By this definition, and because $n/\alpha > 1/Z$, we obtain that $N_{a_{i+1}} \geq (n/\alpha)a_iN_{a_i} > 5^iN_{a_i}$ for each $i < j$. From these inequalities we derive that $N_{a_j} > 5^{\sum_{i=0}^{j-1} i}$. But $N_{a_j} \leq n$, so $\sum_{i=1}^{j-1} i = j(j-1)/2 \leq \log_5 n$. This inequality implies that $j < 1 + \sqrt{2 \log_5 n} < 1 + \sqrt{\lg n}$. By Corollary 2, the height of the BFS tree rooted at v is at most $2(4a_{1+\sqrt{\lg n}} + 1) + 24 \lg n$. Solving the recurrence relation, we obtain that $a_j = O(5^{\sqrt{\lg n}} \lg n)$. By Lemma 2, the price of anarchy is $O(5^{\sqrt{\lg n}} \lg n)$. \square

7. UPPER BOUNDS FOR MAX UNILATERAL GAME

In this section, we introduce and analyze the natural max variation on the (unilateral) network creation game. This problem is motivated by players forming a network with guaranteed worst-case performance subject to budget constraints. We prove that the price of anarchy is at most 2 for $\alpha \geq n$, $O(\min\{4^{\sqrt{\lg n}}, (n/\alpha)^{1/3}\})$ for $2\sqrt{\lg n} \leq \alpha \leq n$, and $O(n^{2/\alpha})$ for $\alpha < 2\sqrt{\lg n}$.

Let D_v be the distance of the furthest vertex from v and $N_k^-(v)$ be the set of vertices whose distance to v is exactly k . Thus the cost incurred by vertex v is $\alpha e_v + D_v$, where e_v is the number of edges that v has bought.

LEMMA 7. Any equilibrium graph G_s has no cycle of length less than $\alpha + 2$.

PROOF. Suppose for contradiction that there is cycle C with $|C| < \alpha + 2$. Let $\{u, v\}$ be an edge of this cycle. Suppose by symmetry that u bought this edge. If vertex u removed the edge, it would decrease its buying cost e_u by α and increase its distance cost D_u by at most $|C| - 2 < \alpha$. Hence it is cost effective for u to remove this edge, contradicting equilibrium. \square

THEOREM 5. For $\alpha \geq n$, the price of anarchy is at most 2.

PROOF. By Lemma 7, the equilibrium graph G_s has no cycle of length at most n , which implies that G is a tree. The cost of s is at most $\alpha(n-1) + \sum_{v \in V(G)} D_v \leq \alpha(n-1) + n(n-1) \leq 2\alpha(n-1)$. On the other hand, the cost of the social optimum is at least $\alpha(n-1)$. Therefore the price of anarchy is at most $\frac{2\alpha(n-1)}{\alpha(n-1)} = 2$. \square

Lemma 7 gives us a lower bound on the *girth* (length of the shortest cycle) of an equilibrium graph. We use the following result of [7] to relate the number of edges to the girth:

LEMMA 8. [7] The number of edges in an n -vertex graph of odd girth g is $O(n^{1+2/(g-1)})$.

LEMMA 9. The number of edges in an equilibrium graph is $O(n^{1+2/\alpha})$.

PROOF. By Lemma 7, the girth of the equilibrium graph G is at least $\alpha + 2$. So by Lemma 8, the number of edges in G is $O(n^{1+2/(\alpha+2-1)})$ for odd values of α , and the number of edges is $O(n^{1+2/(\alpha+1-1)})$ for even values of α . \square

LEMMA 10. For $k \leq D_v/2$, we have $|N_{k+1}^-(v)| \geq k/\alpha$ and $|N_{k+1}(v)| > k^2/(2\alpha)$.

PROOF. If vertex v bought the $|N_{k+1}^-(v)|$ edges connecting v to the vertices in $N_{k+1}^-(v)$, it would decrease D_v by at least $D_v - \max\{D_v - k, k\} = D_v - (D_v - k) = k$. The cost of buying these edges is $|N_{k+1}^-(v)|\alpha$. Because v has not bought these edges, we conclude that this cost is at least k . Hence $|N_{k+1}^-(v)| \geq k/\alpha$. Therefore $|N_{k+1}(v)| \geq \sum_{i=1}^k (i/\alpha) > k^2/(2\alpha)$. \square

LEMMA 11. The diameter of an equilibrium graph G_s , $\text{diam}(G_s)$, is $O((n\alpha^2)^{1/3})$.

PROOF. Set $k = \text{diam}(G_s)/4 - 1$. Let v be some vertex with $D_v = \text{diam}(G_s)$. Similar to the proof of Lemma 6, we select a subset of vertices as *center points* by the following greedy algorithm. First we unmark all vertices in G_s . Then we repeatedly select an unmarked vertex x as a center point, and mark all unmarked vertices whose distances are at most $2k$ from x .

Suppose that we select l vertices x_1, x_2, \dots, x_l as center points. By construction, every vertex in the graph has distance at most $2k$ to some center point. If vertex v bought the l edges $\{v, x_1\}, \{v, x_2\}, \dots, \{v, x_l\}$, it would decrease D_v by at least $D_v - (2k+1) \geq (4k+4) - (2k+1) > 2k$. Buying these edges costs αl . Because v has not bought these edges, we have $\alpha l \geq 2k$. On the other hand, according to the greedy algorithm, the distance between any pair of center points is at least $2k + 1$. Thus we have $N_k(x_i) \cap N_k(x_j) = \emptyset$ for $1 \leq i < j \leq l$. Therefore, $|\bigcup_{i=1}^l N_k(x_i)| = \sum_{i=1}^l |N_k(x_i)|$, which should be less than or equal to n . By Lemma 10, we have $n \geq |\bigcup_{i=1}^l N_k(x_i)| = \sum_{i=1}^l |N_k(x_i)| \geq l(k-1)^2/(2\alpha)$, which means that $l \leq 2n\alpha/(k-1)^2$. Combining with $\alpha l \geq 2k$, we obtain that $2k \leq 2n\alpha^2/(k-1)^2$, i.e., $k(k-1)^2 \leq n\alpha^2$. Therefore $\text{diam}(G_s) = O((n\alpha^2)^{1/3})$. \square

THEOREM 6. For $6 \leq \alpha < n$, the price of anarchy is $O((n/\alpha)^{1/3})$.

PROOF. By Lemma 9, the number of edges in G_s is $O(n^{1+2/\alpha})$. By Lemma 11, $\text{diam}(G_s) = O((n\alpha^2)^{1/3})$. Hence the cost of G_s is at most $\alpha O(n^{1+2/\alpha}) + n O((n\alpha^2)^{1/3})$. On the other hand, the cost of the social optimum is $\Omega(n\alpha)$. Therefore the price of anarchy is $O(n^{2/\alpha} + (n\alpha^2)^{1/3}/\alpha) \leq O(n^{2/\alpha} + (n/\alpha)^{1/3}) = O((n/\alpha)^{1/3})$. \square

LEMMA 12. If $|N_k(u)| > Y$ for every vertex u in an equilibrium graph G_s , then either $D_u \leq 5k$ for some vertex u or $|N_{4k+1}(u)| > Yk/\alpha$ for every vertex u .

PROOF. The proof is similar to the proof of Lemma 6. If there is a vertex u with $D_u \leq 5k$, then the claim is obvious. Otherwise, for every vertex u , we have $D_u > 5k$. Let u be an arbitrary vertex. Let S be the set of vertices whose distance from u is $3k + 1$. We select a subset of S , called *center points*, by the following greedy algorithm. First we unmark all vertices in S . Then we select an unmarked vertex $x \in S$

as a center point, mark all unmarked vertices in S whose distance from x is at most $2k$, and assign these vertices to x .

Suppose that we select l vertices x_1, x_2, \dots, x_l as center points. We prove that $l \geq k/\alpha$. If vertex u bought the l edges $\{u, x_1\}, \{u, x_2\}, \dots, \{u, x_l\}$, it would decrease D_u by at least $D_u - \max\{D_u - k, 3k + 1\} = D_u - (D_u - k) = k$. Because u has not bought these edges, we must have $l\alpha \geq k$.

According to the greedy algorithm, the distance between any pair of center points is more than $2k$; hence $N_k(x_i) \cap N_k(x_j) = \emptyset$ for $i \neq j$. By the hypothesis of the lemma, $|N_k(x_i)| > Y$ for every vertex x_i ; hence $|\bigcup_{i=1}^l N_k(x_i)| = \sum_{i=1}^l |N_k(x_i)| > lY$. For every $i \leq l$, we have $d_s(u, x_i) = 3k + 1$, so vertex u has a path of length at most $4k + 1$ to every vertex whose distance to x_i is at most k . Therefore $|N_{4k+1}(u)| \geq |\bigcup_{i=1}^l N_k(x_i)| > lY \geq Yk/\alpha$. \square

THEOREM 7. *The price of anarchy is $O(4^{\sqrt{\lg n}} + n^{2/\alpha})$.*

PROOF. The proof is similar to the proof of Theorem 4. Let $Z = \alpha$. Consider an arbitrary vertex v . Because any equilibrium graph is connected, $|N_Z| > Z$. By Lemma 12, either $D_v \leq 5k$ for some vertex v or $N_{4k+1} > (k/\alpha)N_k$ for every vertex v . Define the numbers a_0, a_1, \dots using the recurrence relation $a_i = 4a_{i-1} + 1$ with $a_0 = Z$. By induction, $a_i \geq Z4^i$. Suppose that j is the least number for which $D_v \leq 5a_j$ for some vertex v . By this definition, $N_{a_{i+1}} \geq (a_i/\alpha)N_{a_i} \geq 4^i N_{a_i}$ for each $i < j$. From these inequalities we derive that $N_{a_j} > 4^{\sum_{i=0}^{j-1} i}$. But $|N_{a_j}(v)| \leq n$, so $\sum_{i=1}^{j-1} i = j(j-1)/2 \leq \log_4 n$. This inequality implies that $j \leq 1 + \sqrt{2 \log_4 n} = 1 + \sqrt{\lg n}$. Also, $D_v \leq 5a_j$. Solving the recurrence relation, $a_j = O(4^j \alpha)$. Thus D_v and therefore $\text{diam}(G_s)$ are $O(4^{\sqrt{\lg n}} \alpha)$.

On the other hand, by Lemma 9, the number of edges in the graph is $O(n^{1+2/\alpha})$. Hence the cost of the graph is at most $n \text{diam}(G_s) + \alpha O(n^{1+2/\alpha}) = O(n\alpha 4^{\sqrt{\lg n}} + \alpha n^{1+2/\alpha})$. The cost of the social optimum is at least $\alpha(n-1) = \Omega(n\alpha)$. Therefore the price of anarchy is $O(4^{\sqrt{\lg n}} + n^{2/\alpha})$. \square

We conclude the following corollaries from Theorem 7:

COROLLARY 3. *For $\alpha > \sqrt{\lg n}$, the price of anarchy is $O(4^{\sqrt{\lg n}})$.*

COROLLARY 4. *For $\alpha \leq \sqrt{\lg n}$, the price of anarchy is $O(n^{2/\alpha})$.*

8. TIGHT LOWER BOUNDS FOR SUM BILATERAL GAME

In this section, we prove tight lower bounds on the sum version of the bilateral network creation game. Although not stated explicitly, [5, Proposition 4] establishes an upper bound of $O(\frac{n\sqrt{\alpha}}{\alpha+n})$. For $\alpha < n$, this upper bound is $O(\sqrt{\alpha})$; for $\alpha > n$, this upper bound is $O(n/\sqrt{\alpha})$. We prove matching lower bounds using the following lemma:

LEMMA 13. *There is a tree $T_{d,k}$ with $n = 1 + dk$ vertices and the total cost greater than $2\alpha(n-1) + C(k, 2)d^2(d+1) = 2\alpha(n-1) + k(k-1)d^2(d+1)$, for $\alpha > 2d^2$.*

PROOF. We construct the tree $T_{d,k}$ as follows. Put a vertex r as root in the tree. Attach k paths P_1, P_2, \dots, P_k each of which of length d to r . So there are k paths with d edges

which all of them have r as one of their endpoints. Note that there are kd edges and $n = 1 + kd$ vertices in this tree.

Because all edges are cut edges, there is no edge $\{u, v\}$ that u and v want to remove. On the other hand, it is sufficient to prove that no edge will be added, to conclude that $T_{d,k}$ is an equilibrium graph. For sake of contradiction, suppose there is a pair of vertices such as a and b such that both of them want to build the edge $\{a, b\}$ and pay α for the edge. Suppose that $a \in P_i$ and $b \in P_j$, note that i may be equal to j which means that a and b are from one path. Let d_a and d_b be the length of the paths from r to a and b in $T_{d,k}$ respectively. One of the d_a and d_b might be zero which means that one of the a and b is the root r . Without loss of generality, assume that $d_a \leq d_b$. The length of the shortest path from a to the vertices out of path P_j would not decrease with this edge. On the other hand its distance to vertices in P_j is decreased at most $|P_i| + |P_j| = 2d$. Because there are d vertices in P_j , vertex a has decreased its sum of distances to the other $n-1$ vertices at most $d(2d) = 2d^2$. But a paid α to buy this edge. So $\alpha \leq 2d^2$ which is a contradiction. Therefore $T_{d,k}$ is an equilibrium graph. The cost of $T_{d,k}$ is $\alpha(n-1) + \sum_{u,v \in V(T)} d_s(u, v)$. The sum of the distances between vertices in different paths is obviously less than the sum of distances over all pairs of vertices. The distance between two vertices from two different paths is equal to the sum of their distances to r . So the cost of the tree is at least $2\alpha(n-1) + \sum_{i \neq j} \sum_{a=1}^d \sum_{b=1}^d a + b = 2\alpha(n-1) + k(k-1)d^2(d+1)$. \square

THEOREM 8. *The price of anarchy is at least $\Omega(\sqrt{\alpha})$ when $\alpha < n$.*

PROOF. By Lemma 13 and setting $d = \sqrt{\alpha/2} - 1$, we reach an equilibrium graph with cost at least $d^2(d+1)k(k-1) = \Omega(n^2d)$. The social optimum (a star) has cost at most $2\alpha(n-1) + 2n^2 = O(n^2)$. Therefore the price of anarchy is at least $\Omega(d) = \Omega(\sqrt{\alpha})$. \square

THEOREM 9. *The price of anarchy is at least $\Omega(n/\sqrt{\alpha})$ when $\alpha \geq n$.*

PROOF. Again by Lemma 13 and setting $d = \sqrt{\alpha/2} - 1$, we reach an equilibrium graph with cost at least $d^2(d+1)k(k-1) = \Omega(n^2d)$. The social optimum (a star) has cost at most $2\alpha(n-1) + 2n^2 = O(\alpha n)$. Therefore the price of anarchy is at least $\Omega(nd/\alpha) = \Omega(n/\sqrt{\alpha})$. \square

9. TIGHT BOUNDS FOR MAX BILATERAL GAME

In this section, we prove tight bounds on the max version of the bilateral network creation game.

THEOREM 10. *The price of anarchy is at least $\Omega(\frac{n}{\alpha+1})$ for any α .*

PROOF. Consider tree $T_{d,3}$ as it is defined in Lemma 13 which has $n = 1 + 3d$ vertices. Again because all edges are cut edges, there is no edge $\{u, v\}$ that u and v want to remove. On the other hand, suppose there is a pair of vertices such as a and b such that both of them want to build the edge $\{a, b\}$ and pay α for the edge. Assume that $a \in P_i$ and $b \in P_j$, note that i may be equal to j which means that a and b are from one path. Let d_a and d_b be the length of the paths from r to a and b in $T_{d,3}$ respectively.

One of the d_a and d_b might be zero which means that one of the a and b is the root r . Without loss of generality, assume that $d_a \leq d_b$. After adding this edge, the height of the BFS tree rooted at a remains $d_a + d$ which has not changed. So vertex a has no interest to buy this edge. Therefore $T_{d,3}$ is an equilibrium graph.

The cost of $T_{d,3}$ is at least $\Omega(2\alpha(n-1) + nd) = \Omega(\alpha n + n^2) = \Omega(n^2)$. Because the cost of the social optimum is $O(\alpha n + n)$, the price of anarchy is at least $\Omega(\frac{n}{\alpha+1})$. \square

THEOREM 11. *The price of anarchy is $O(\frac{n}{\alpha+1})$ for any $\alpha \leq n$, and at most 2 for $\alpha > n$.*

PROOF. Because players can force the removal of edges in the bilateral version, Lemma 7 still holds by the same proof. Thus we can apply Theorem 5 to obtain the desired upper bound of 2 in the case $\alpha > n$. When $\alpha \leq n$, we can apply Lemma 9 to derive that no cycle has length less than $\alpha + 2$ in an equilibrium graph. Because the number of edges in the graph is also at most $n(n-1)/2$, there are $O(\min\{n^2, n^{1+2/\alpha}\})$ edges in the equilibrium graph. On the other hand, the height of the BFS tree rooted at any vertex of the graph is at most n . So the cost is $O(\min\{n^2, n^{1+2/\alpha}\}\alpha + n^2)$. Knowing the fact that the social optimum cost is at least $\Omega(\alpha n + n)$, the price of anarchy is $\frac{O(n^{1+2/\alpha}\alpha + n^2)}{\Omega(\alpha n + n)} = O(n^{2/\alpha} + \frac{n}{\alpha+1}) = O(\frac{n}{\alpha+1})$ for $2 < \alpha \leq n$ and $\frac{O(n^2\alpha + n^2)}{\Omega(\alpha n + n)} = O(n) = O(\frac{n}{\alpha+1})$ for $\alpha \leq 2$. Therefore, for any $\alpha \leq n$, the price of anarchy is $O(\frac{n}{\alpha+1})$. \square

10. CONCLUSION

In this paper, we have bounded the price of anarchy in four different network creation games. We have significantly improved the bounds for the sum unilateral game, introduced the new max game, and completely resolved the bilateral games. We conjecture that the correct bound on the price of anarchy for the sum unilateral game is $\Theta(1)$. For the max unilateral game, a general constant bound does not seem impossible; in any case, it would be interesting to determine the optimal bound.

One interesting generalization of all of these games is when only some links can possibly be created (because of physical limitations, for example). More precisely, we are given a (connected) graph of the allowable edges, and the players correspond to nodes in this graph. In this case, the socially optimal strategy is no longer simply a clique or a star, and it is not even clear whether it can be computed in polynomial time. Thus the price of stability (the *minimum* cost of a Nash equilibrium divided by the social optimum) also becomes of interest.

11. REFERENCES

- [1] Susanne Albers, Stefan Eilts, Eyal Even-Dar, Yishay Mansour, and Liam Roditty. On Nash equilibria for a network creation game. In *Proceedings of the 17th Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 89–98, Miami, Florida, 2006.
- [2] Elliot Anshelevich, Anirban Dasgupta, Jon Kleinberg, Eva Tardos, Tom Wexler, and Tim Roughgarden. The price of stability for network design with fair cost allocation. In *Proceedings of the 45th Annual IEEE Symposium on Foundations of Computer Science*, pages 295–304, 2004.
- [3] Elliot Anshelevich, Anirban Dasgupta, Eva Tardos, and Tom Wexler. Near-optimal network design with selfish agents. In *Proceedings of the 35th Annual ACM Symposium on Theory of Computing*, pages 511–520, San Diego, California, 2003.
- [4] Byung-Gon Chun, Rodrigo Fonseca, Ion Stoica, and John Kubiatowicz. Characterizing selfishly constructed overlay routing networks. In *Proceedings of the 23rd Annual Joint Conference of the IEEE Computer and Communications Societies*, Hong Kong, China, March 2004.
- [5] Jacomo Corbo and David Parkes. The price of selfish behavior in bilateral network formation. In *Proceedings of the 24th Annual ACM Symposium on Principles of Distributed Computing*, pages 99–107, Las Vegas, Nevada, 2005.
- [6] Artur Czumaj and Berthold Vöcking. Tight bounds for worst-case equilibria. In *Proceedings of the 13th Annual ACM-SIAM symposium on Discrete Algorithms*, pages 413–420, San Francisco, California, 2002.
- [7] R. D. Dutton and R. C. Brigham. Edges in graphs with large girth. *Graphs and Combinatorics*, 7(4):315–321, December 1991.
- [8] Alex Fabrikant, Ankur Luthra, Elitza Maneva, Christos H. Papadimitriou, and Scott Shenker. On a network creation game. In *Proceedings of the 22nd Annual Symposium on Principles of Distributed Computing*, pages 347–351, Boston, Massachusetts, 2003.
- [9] Matthew O. Jackson. A survey of models of network formation: Stability and efficiency. In Gabrielle Demange and Myrna Wooders, editors, *Group Formation in Economics; Networks, Clubs and Coalitions*. Cambridge University Press, 2003.
- [10] Elias Koutsoupias and Christos H. Papadimitriou. Worst-case equilibria. In *Proceedings of the 16th Annual Symposium on Theoretical Aspects of Computer Science*, volume 1563 of *Lecture Notes in Computer Science*, pages 404–413, Trier, Germany, March 1999.
- [11] Tom Leighton and Satish Rao. Multicommodity max-flow min-cut theorems and their use in designing approximation algorithms. *Journal of the ACM*, 46(6):787–832, 1999.
- [12] Henry Lin. On the price of anarchy of a network creation game. Class final project, December 2003.
- [13] Christos Papadimitriou. Algorithms, games, and the internet. In *Proceedings of the 33rd Annual ACM Symposium on Theory of Computing*, pages 749–753, Hersonissos, Greece, 2001.
- [14] Tim Roughgarden. The price of anarchy is independent of the network topology. In *Proceedings of the 34th Annual ACM Symposium on Theory of Computing*, pages 428–437, Montréal, Canada, 2002.
- [15] Tim Roughgarden. *Selfish Routing*. PhD thesis, Cornell University, 2002. Published as *Selfish Routing and the Price of Anarchy*, MIT Press, 2005.